

# On polynomial solutions of the Lamé and Stokes systems

Hiroya ITO\*

The University of Electro-Communications

## 1 Introduction

The Laplacian  $\Delta$  is one of the most important differential operators in Mathematics. Solutions of the Laplace equation  $\Delta u = 0$  are called harmonic functions, which play significant roles in many subjects of mathematical research fields. It is well known that harmonic polynomials in  $n$  variables are well classified. Moreover, the restriction of nonzero elements of  $\mathcal{H}_m$  to the unit sphere  $\mathbb{S}^{n-1}$ , called spherical harmonics of degree  $m$ , become eigenfunctions of the Laplace-Beltrami operator  $-\Delta_{\mathbb{S}^{n-1}}$  on  $\mathbb{S}^{n-1}$  with the common eigenvalue  $m(m+n-2)$ , and this restriction is a linear isomorphism between  $\mathcal{H}_m$  and the space of spherical harmonics of degree  $m$ . Furthermore, we can think that all the harmonic polynomials in  $\mathbb{R}^n$  (or all the spherical harmonics) generate most function spaces on  $\mathbb{S}^{n-1}$  (see Theorem 1 below). We shall study that such beautiful theory can be partially generalized to vector-valued elliptic systems.

Fixing  $n$  variables  $x_1, \dots, x_n$  with  $n \geq 2$ , for each  $m \in \mathbb{N}_0$ , we denote by  $\mathcal{P}_m$  the vector space of all the homogeneous polynomials of degree  $m$  in  $x = (x_1, \dots, x_n)$ , and by  $\mathcal{H}_m$  its subspace consisting of those in  $\mathcal{P}_m$  which are harmonic. Moreover,  $\mathring{\mathcal{H}}_m$  denotes the vector space of all the functions on  $\mathbb{S}^{n-1}$  obtained by restricting each element of  $\mathcal{H}_m$  to  $\mathbb{S}^{n-1}$ ; each element of  $\mathring{\mathcal{H}}_m$  is called a spherical harmonic of degree  $m$ :

$$\mathcal{H}_m = \{u \in \mathcal{P}_m \mid \Delta u = 0\}, \quad \mathring{\mathcal{H}}_m = \{u|_{\mathbb{S}^{n-1}} \mid u \in \mathcal{H}_m\}.$$

Then, the dimension  $d_m$  of  $\mathcal{P}_m$  is given by  $d_m = \binom{m+n-1}{n-1}$  and the restriction map  $\mathcal{H}_m \ni u \mapsto u|_{\mathbb{S}^{n-1}} \in \mathring{\mathcal{H}}_m$  is, due to the homogeneity of elements of  $\mathcal{H}_m$ , a linear isomorphism:  $\mathcal{H}_m \cong \mathring{\mathcal{H}}_m$ . Fundamental properties of spherical harmonics on  $\mathbb{S}^{n-1}$  are described in the following theorem (see, e.g., Chapter 2 of Shimakura [4], Chapter 3 of Simon [5], Nomura [2]).

**Theorem 1.** *The space  $\mathring{\mathcal{H}}_m$  ( $m \in \mathbb{N}_0$ ) has the following properties.*

(i) *The dimension of  $\mathring{\mathcal{H}}_m$  is given by  $\dim \mathring{\mathcal{H}}_m = d_m - d_{m-2}$ , where  $d_{-1} = d_{-2} = 0$ .*

(ii)  *$L^2(\mathbb{S}^{n-1}) = \bigoplus_{m=0}^{\infty} \mathring{\mathcal{H}}_m$  in the sense that*

$$\mathring{\mathcal{H}}_\ell \perp \mathring{\mathcal{H}}_m \text{ (} \ell \neq m \text{) in } L^2(\mathbb{S}^{n-1}) \quad \text{and} \quad \overline{\text{span}(\bigcup_{m=0}^{\infty} \mathring{\mathcal{H}}_m)}^{L^2} = L^2(\mathbb{S}^{n-1}).$$

In the present paper, we consider the homogeneous equation of the Lamé system

$$\mathcal{L}\mathbf{u} := \mu\Delta\mathbf{u} + (\lambda + \mu)\nabla(\text{div}\mathbf{u}) = \mathbf{0} \quad \text{in } \mathbb{R}^n \tag{1}$$

for  $n$ -vector valued functions (vector fields)  $\mathbf{u}$ , where  $\lambda$  and  $\mu$  are elasticity constants. We study the structure of the restriction of polynomial solutions of (1) to  $\mathbb{S}^{n-1}$  (or analogues of spherical harmonics

---

\*e-mail: ito-hiroya@uec.ac.jp

for (1)). Here, the operator  $\mathcal{L}$  of (1) appears in linear theory of isotropic elasticity and the constants  $\lambda$  and  $\mu$  are assumed to satisfy

$$\mu(\lambda + 2\mu) > 0, \quad \gamma := \frac{\lambda + \mu}{\lambda + 3\mu} \in (-1, 1). \quad (2)$$

The symbol of  $\mathcal{L}$  is

$$L(\xi) = \mu|\xi|^2 I + (\lambda + \mu)\boldsymbol{\xi} \otimes \boldsymbol{\xi}, \quad (3)$$

whose eigenvalues are given by  $\mu|\xi|^2$  (multiplicity  $n - 1$ ) and  $(\lambda + 2\mu)|\xi|^2$  (simple), where we write  $\boldsymbol{\xi}$  in boldface in order to clarify that  $\boldsymbol{\xi}$  is a column vector. Assumption (2) implies that  $\mathcal{L} = L(\partial)$  is a strongly elliptic system. In a similar way we also deal with polynomial solutions of the homogeneous equations of the Stokes system.

## 2 Orthogonally invariant partial differential operators for vector fields

Let  $P(\partial)$  be a partial differential operator with constant coefficients for scalar fields  $u$  on  $\mathbb{R}^n$ . It is well-known that  $P(\partial)$  is invariant under the special orthogonal group  $\text{SO}(n)$  if and only if  $P(\partial)$  is in the form  $P(\partial)u = f(\Delta)u$  for some polynomial  $f(t)$ . How about the case  $P(\partial)$  is for vector fields  $\mathbf{u}$  on  $\mathbb{R}^n$ ? The following theorem In the case  $P(\partial)$  for scalar functions  $u$ , it is well-known that  $P(\partial)$  is invariant under  $\text{SO}(n)$  if and only if it is in the form  $P(\partial)u = f(\Delta)u$ . The following theorem shows, in a sense, the necessity of considering the Lamé system.

**Theorem 2.** *A partial differential operator  $P(\partial)$  with constant coefficients for vector fields  $\mathbf{u}$  on  $\mathbb{R}^n$  is invariant under the orthogonal group  $\text{O}(n)$  if and only if  $P(\partial)$  is in the form*

$$P(\partial)\mathbf{u} = f(\Delta)\mathbf{u} + g(\Delta)\nabla(\text{div } \mathbf{u}).$$

for some polynomials  $f(t)$  and  $g(t)$ .

Even if we restrict  $\text{O}(n)$  to  $\text{SO}(n)$  in Theorem 2, then the conclusion is valid for  $n \geq 4$ , but not for  $n = 2, 3$ , in which  $P(\partial)\mathbf{u}$  may contain additional terms, for example  $h(\Delta)\text{rot } \mathbf{u}$  if  $n = 3$ .

## 3 $L$ -harmonic vector fields and $L$ -harmonics

Denote by  $\mathcal{P}_m$  the vector space of all  $n$ -vector homogeneous polynomials in  $x = (x_1, \dots, x_n)$  of degree  $m$ . We define subspaces  $\mathcal{H}_m$  and  $\mathcal{H}_m^L$  of  $\mathcal{P}_m$  by

$$\mathcal{H}_m = \{\mathbf{u} \in \mathcal{P}_m \mid \Delta \mathbf{u} = \mathbf{0}\}, \quad \mathcal{H}_m^L = \{\mathbf{u} \in \mathcal{P}_m \mid \mathcal{L} \mathbf{u} = \mathbf{0}\}.$$

Elements of  $\mathcal{H}_m^L$  are called  *$L$ -harmonic polynomials* of degree  $m$ .

Vector functions on  $\mathbb{S}^{n-1}$  obtained by restricting  $L$ -harmonic polynomials are called *spherical  $L$ -harmonics*. We represent the vector spaces of such vector functions (vector fields) as

$$\mathring{\mathcal{H}}_m = \{\mathbf{u}|_{\mathbb{S}^{n-1}} \mid \mathbf{u} \in \mathcal{H}_m\}, \quad \mathring{\mathcal{H}}_m^L = \{\mathbf{u}|_{\mathbb{S}^{n-1}} \mid \mathbf{u} \in \mathcal{H}_m^L\}.$$

Corresponding to Theorem 1 for spherical harmonics, the following theorem for spherical  $L$ -harmonics has been established through joint research with Prof. Honda and Prof. Jimbo.

**Theorem 3** ([1]). *The space  $\mathring{\mathcal{H}}_m^L$  ( $m \in \mathbb{N}_0$ ) has the following properties.*

- (i) *The dimension of  $\mathring{\mathcal{H}}_m^L$  is given by  $\dim \mathring{\mathcal{H}}_m^L = \dim \mathcal{H}_m^L = n(d_m - d_{m-2})$ , where  $d_{-1} = d_{-2} = 0$ .*
- (ii) *For each  $m \in \mathbb{N}$ , the sum  $\mathring{\mathcal{H}}_0^L + \mathring{\mathcal{H}}_1^L + \dots + \mathring{\mathcal{H}}_m^L$  is a direct sum.*
- (iii) *The linear span of  $\bigcup_{m=0}^{\infty} \mathring{\mathcal{H}}_m^L$  is dense in  $\mathbf{L}^2(\mathbb{S}^{n-1})$  with the  $L^2$ -norm.*

## 4 Case $n = 2$

In this section we consider the case  $n = 2$ . The spaces  $\mathcal{H}_m, \mathcal{H}_m^L, \mathring{\mathcal{H}}_m, \mathring{\mathcal{H}}_m^L$  are defined not only for nonnegative integer  $m$  but also negative integer  $m$ . For example,  $\mathbf{u} \in \mathring{\mathcal{H}}_m^L$  for  $m < 0$  implies that  $\mathbf{u}$  is a vector field solution of (1) in  $\mathbb{R}^2 \setminus \{0\}$  which is homogeneous in  $x = (x_1, x_2)$  of degree  $m$ .

Let  $\mathbf{u} = (u_1, u_2)$  be a real vector field solution of  $\mathcal{L}\mathbf{u} = \mathbf{0}$  in  $\mathbb{R}^2 \setminus \{0\}$ . Then the complex function  $U(z) := u_1(x_1, x_2) + iu_2(x_1, x_2)$  ( $z := x_1 + ix_2$ ) satisfies

$$\frac{\partial}{\partial \bar{z}} \left( \frac{\partial U}{\partial z} + \gamma \frac{\partial \bar{U}}{\partial z} \right) = 0 \quad \text{in } \mathbb{C} \setminus \{0\},$$

which is solved as

$$U = \varphi(z) - \gamma z \overline{\varphi'(z)} + \overline{\psi(z)} + 2c \log|z| - \gamma \bar{c} \left( \frac{z}{|z|} \right)^2$$

where  $\varphi(z), \psi(z)$  are holomorphic functions in  $\mathbb{C} \setminus \{0\}$ , and  $c \in \mathbb{C}$  is a constant ([3]). Using this fact, we have the following assertions.

**Theorem 4.** *The spaces  $\mathcal{H}_m^L$  and  $\mathring{\mathcal{H}}_m^L$  have the following bases.*

(i)  $\mathcal{H}_0^L = \mathbb{R}^2$ . For  $m \neq 0$ , the space  $\mathcal{H}_m^L$  has a basis

$$\left\{ \begin{bmatrix} \operatorname{Re}[z^m - \gamma m z \overline{z^{m-1}}] \\ \operatorname{Im}[z^m - \gamma m z \overline{z^{m-1}}] \end{bmatrix}, \begin{bmatrix} -\operatorname{Im}[z^m + \gamma m z \overline{z^{m-1}}] \\ \operatorname{Re}[z^m + \gamma m z \overline{z^{m-1}}] \end{bmatrix}, \begin{bmatrix} \operatorname{Re}[z^m] \\ -\operatorname{Im}[z^m] \end{bmatrix}, \begin{bmatrix} \operatorname{Im}[z^m] \\ \operatorname{Re}[z^m] \end{bmatrix} \right\}.$$

(ii)  $\mathring{\mathcal{H}}_0^L = \mathbb{R}^2$ . For  $m \neq 0$ , the space  $\mathring{\mathcal{H}}_m^L$  has a basis

$$\left\{ \begin{bmatrix} \cos m\theta - \gamma m \cos(m-2)\theta \\ \sin m\theta + \gamma m \sin(m-2)\theta \end{bmatrix}, \begin{bmatrix} -\sin m\theta + \gamma m \sin(m-2)\theta \\ \cos m\theta + \gamma m \cos(m-2)\theta \end{bmatrix}, \begin{bmatrix} \cos m\theta \\ -\sin m\theta \end{bmatrix}, \begin{bmatrix} \sin m\theta \\ \cos m\theta \end{bmatrix} \right\}.$$

**Corollary 5.**

(i)  $\mathring{\mathcal{H}}_0^L + \mathring{\mathcal{H}}_1^L + \cdots + \mathring{\mathcal{H}}_m^L = \mathring{\mathcal{H}}_0 + \mathring{\mathcal{H}}_1 + \cdots + \mathring{\mathcal{H}}_m$  for  $m \geq 1$ .

(ii) The sum  $\mathring{\mathcal{H}}_0^L + \mathring{\mathcal{H}}_{-1}^L + \cdots + \mathring{\mathcal{H}}_{-m}^L$  ( $m \geq 1$ ) is a direct sum, and satisfies

$$\mathring{\mathcal{H}}_0 + \mathring{\mathcal{H}}_1 + \cdots + \mathring{\mathcal{H}}_{m-2} \subset \mathring{\mathcal{H}}_0^L + \mathring{\mathcal{H}}_{-1}^L + \cdots + \mathring{\mathcal{H}}_{-m}^L \subset \mathring{\mathcal{H}}_0 + \mathring{\mathcal{H}}_1 + \cdots + \mathring{\mathcal{H}}_{m+2} \quad \text{for } m \geq 2.$$

## References

- [1] N. Honda, H. Ito and S. Jimbo, Spherical vector functions associated with the Lamé and Stokes systems, *preprint*.
- [2] T. Nomura, Spherical Harmonics and Group Representations (*Japanese*), NipponHyoronsha (2018).
- [3] N. I. Muskhelishvili, Some Basic Problems of the Mathematical Theory of Elasticity, 2nd ed., Noordhoff N.V., 1953.
- [4] N. Shimakura, Partial Differential Operators of Elliptic Type, AMS, Providence, RI, 1992.
- [5] B. Simon, Harmonic analysis, A Comprehensive Course in Analysis, Part 3. AMS, Providence, RI, 2015.